

PRELIMINARY RESULTS FROM THE ESO SLICE PROJECT (ESP)

G. Vettolani and E. Zucca

Istituto di Radioastronomia del CNR, Bologna, Italy

A. Cappi, R. Merighi, M. Mignoli, G. Stirpe and G. Zamorani

Osservatorio Astronomico di Bologna, Bologna, Italy

C. Collins and H. MacGillivray

Royal Observatory Edinburgh, Edinburgh, United Kingdom

C. Balkowski, J. Alimi, A. Blanchard, V. Cayatte,

P. Felenbok, S. Maurogordato and D. Proust

DAEC, Observatoire de Paris–Meudon, Meudon, France

G. Chincarini and L. Guzzo

Osservatorio Astronomico di Brera, Milano, Italy

D. Maccagni

Istituto di Fisica Cosmica e Tecnologie Relative, Milano, Italy

R. Scaramella

Osservatorio Astronomico di Roma, Monteporzio Catone, Italy

M. Ramella

Osservatorio Astronomico di Trieste, Trieste, Italy

Abstract

We present the first results of a galaxy redshift survey, ESO Slice Project (ESP), we are accomplishing as an ESO Key–Project over about 40 square degrees in a region near the South Galactic Pole. The limiting magnitude is $b_J = 19.4$. Up to now $\sim 85\%$ of the observations has been completed and $\sim 65\%$ of the data has been reduced providing ~ 2000 galaxy redshifts.

We present some preliminary results concerning the large scale galaxy distribution, the luminosity function and the properties of galaxies with spectra showing prominent emission lines.

1. Introduction

In September 1991 we started a galaxy redshift survey over a strip of $22^\circ \times 1^\circ$ (plus a near–by area of $5^\circ \times 1^\circ$, five degrees west of the strip) in the South Galactic Pole region. The right ascension limits are $22^h 30^m$ and $01^h 20^m$, at a

Fig.1: The survey plan: each circle corresponds to one OPTOPUS field. The spectra of galaxies inside the fields shown as filled circles have been already reduced.

mean declination of -40° (1950). We have filled this area with a regular grid of circular fields with a diameter of 32 arcminutes (see Figure 1). This size corresponds to the field of view of the multifiber spectrograph (OPTOPUS) we use at the ESO $3.6m$ telescope.

The limiting magnitude of the survey is $b_J \leq 19.4$. We have selected the target objects from the Edinburgh–Duhram Southern Sky Galaxy Catalogue (Heydon–Dumbleton et al. 1989), which has been obtained from COSMOS scans of 60 plates in the SGP region. The catalogue has a 95% completeness at $b_J \leq 20.5$, a photometric accuracy of ~ 0.03 magnitudes and an estimated stellar contamination $\leq 10\%$.

The total number of objects we have observed is of the order of 4000. We have obtained spectra for galaxies in all the fields and have reduced the data for 70 of them (shown as filled circles in Figure 1), obtaining 1924 galaxy redshifts.

At $z \simeq 0.1$, which corresponds to the peak of the selection function of the survey, the linear dimensions of the full strip are of the order of $110 \times 5 \text{ } h^{-1} \text{ Mpc}$.

The main goals of our project can be summarized as follows:

- 1) Determination of the galaxy luminosity function (estimating both its shape and normalization) in a volume with dimensions large enough to average over the large scale inhomogeneities.
- 2) Study of the statistics of emission line galaxies (both normal and active) in a large, unbiased sample of galaxies.
- 3) Measure of the size distribution of inhomogeneities in the galaxy distribution over a large volume.

2. Observations and Data Reduction

The observations have been obtained with the multifiber spectrograph OPTOPUS at the Cassegrain focus of the ESO $3.6m$ telescope at La Silla. OPTO-

PUS has 50 fibers which are manually plugged into holes drilled in aluminum plates. The diameter on the sky of the fibers is 2.4 arcseconds . At $b_J \leq 19.4$ the median number of galaxies per OPTOPUS field is 34 and 85% of the fields has less than 45 galaxies. Five fibers are dedicated to the sky observations and the remaining 45 fibers are used to observe galaxies. When the number of galaxies in the field exceeds the number of available fibers, the galaxies to be observed are chosen at random from the galaxy list. However, the overdense fields have been often observed twice to ensure a completeness of the order of 80% or larger.

Under optimal conditions we have observed up to 6 or 7 fields per night. The exposure time for each field is one hour, split into two 30 minute exposures in order to ease the cosmic rays removal. Comparison spectra and white lamp flats have been taken at the beginning and end of each exposure. The spectra cover the wavelength range from $3730 \text{ \AA}$ to $6050 \text{ \AA}$, with an effective resolution of about $3.5 \text{ \AA}$.

Spectra are flat-fielded, extracted and wavelength calibrated. Then the relative transmission of each fiber is computed by normalizing each spectrum through the flux of the OI $\lambda 5577$ and Na $\lambda 5891$ sky lines.

After subtraction of the average sky from the galaxy plus sky spectra (the accuracy of the sky subtraction is in the range 2–5%), the redshifts are measured by cross-correlating the spectra with a set of template stars observed by us with the same instrument. Redshift measurements obtained from the cross-correlation are not considered reliable when the value of the R parameter (Tonry & Davis 1979) is $R \leq 3$, unless the measurement is confirmed by emission lines at the same redshift. The redshifts from emission lines (when present) are also measured, as well as the equivalent width of the most common emission lines. The median error in velocity for the total sample is $\sim 50 \text{ km/s}$.

Figure 2 shows the wedge diagram of the 1924 galaxies with measured redshift. About 80% of the fields observed up to now have a redshift completeness greater than 60%. The less complete fields have been reobserved, but the new data have not been reduced yet.

3. Large Scale Properties

The histogram in Figure 3a shows the distribution in comoving distance ($q_0 = 0.5$) of the 1924 galaxies measured so far. The outstanding peak at $D \simeq 300 h^{-1} \text{ Mpc}$ is not due to a single galaxy cluster but to a structure which extends over almost all the fields (see Figure 2). The distance of this peak is almost coincident with the maximum of our selection function. Other

Fig.2: Wedge diagram in the region $22^h30^m < \alpha < 1^h30^m$ and $-41^\circ15' < \delta < -39^\circ45'$.

peaks seen in Figure 3a at smaller and larger distance are due to less prominent structures, but with a similar contrast with respect to a uniform distribution. These structures are clearly seen in Figure 4, which shows a better representation of the large scale distribution of galaxies in this survey. This figure has been produced by giving to each galaxy in each field a weight proportional to the selection function (derived from the luminosity function described in the next paragraph) and to the incompleteness of the field it belongs to. This procedure implicitly assumes that the galaxies we did not observe follow the same distribution in depth as the observed galaxies. This assumption is justified by the fact that, in overdense fields, the galaxies we observed were selected at random and that the apparent magnitude distributions of the galaxies with and without redshift are fully compatible. Data were subsequently smoothed with a gaussian window of $6 h^{-1} Mpc$.

A surprisingly large fraction of galaxies ($\sim 40\%$) shows the presence of one or more emission lines (OII $\lambda 3727$, $H\beta$, OIII $\lambda 4959$ and $\lambda 5007$). These objects can be either spiral galaxies, where lines originate mostly from HII regions in the disks, or early-type galaxies undergoing a significant burst of star formation.

The distribution in distance of the galaxies with emission lines is shown as a dashed histogram in Figure 3b. It is clear from the figure and also confirmed by

Fig.3: a) Galaxy distribution in comoving distance ($q_0 = 0.5$). The solid line shows the expectation resulting from a uniform distribution of the galaxies in the sample. Vertical lines show the position of the BEKS peaks (see text).
b) Galaxy distribution in comoving distance; the hatched histogram displays only galaxies showing emission lines.

statistical tests, that the large scale distribution of galaxies with emission lines is different from that of galaxies without emission lines: the observed peaks in the galaxy distribution are much less apparent in galaxies with emission lines. This suggests that either spiral galaxies are less frequent in the densest regions, thus confirming a large scale validity of the well known morphology–density relation, or starbursts phenomena occur preferentially in low density environments or both.

After completion of the survey we will be able to explore further this point on the basis of quantitative estimates of densities and a clearer definition of the structures beyond pure visual impression.

Fig.4: Wedge diagram showing the galaxy distribution in the first row of OPTOPUS fields, smoothed with a gaussian window function, taking into account both the incompleteness and the selection function of the sample. The X and Y scales are in $h^{-1} Mpc$ (comoving). Note that East and West are flipped with respect to the previous wedge diagram.

The vertical lines in Figure 3a show the location of the regularly spaced density enhancements as Broadhurst et al. (1990) found in their deep pencil beam survey in the South Galactic Pole region. The two main peaks in our redshift distribution (at comoving distances of ~ 170 and $\sim 300 h^{-1} Mpc$) are in reasonably good agreement with the Broadhurst et al. (1990) peaks and may well be part of the same structures (walls) orthogonal to the line of sight.

Under this hypothesis, since the Broadhurst et al. (1988) survey region is located $\sim 10^\circ$ north of the eastern corner of the present survey, the structure at $z \simeq 0.1$ would have minimum linear dimensions of the order of $110 \times 50 h^{-1} Mpc$, comparable with the Great Wall (Geller & Huchra 1989).

4. The Luminosity Function

The well controlled selection of our sample and the large number of redshifts

already obtained allow us to estimate for the first time the shape parameters of the galaxy luminosity function at magnitudes as faint as $b_J = 19.4$.

Since our database was selected in the blue–green, K–corrections are needed to compute the luminosity function even for the moderate redshifts sampled by our galaxies ($z \leq 0.2$). The functional forms of the K–correction as a function of redshift have been taken from Shanks et al. (1984). Application of the K–correction to each individual galaxy obviously requires the knowledge of its morphological type. In our case the determination of the morphological types by visual inspection of the plates is difficult since most of our galaxies are fainter than 17^{th} magnitude. To overcome this problem, we have adopted the following statistical approach. First, we have assumed that the percentages of late– and early–type galaxies in our sample are approximately the same as those observed in brighter and nearer samples, *i.e.* $\sim 70\%$ and $\sim 30\%$ respectively (Shanks et al. 1984). Then, we have applied the K–correction appropriate for a late–type galaxy to all galaxies showing emission lines ($\sim 40\%$). Finally, we have randomly assigned a morphological type to the remaining galaxies ($\sim 60\%$) in such a way as to obtain the assumed ratio of late and early types in the total sample. A number of tests performed by varying this ratio show that the final results are only mildly dependent on the adopted value.

Following the STY method of Sandage et al. (1979), we have then derived the parameters of the Schechter functional form of the luminosity function, through a maximum likelihood technique. The best fit parameters are $\alpha = -1.13$ and $M_{b_J}^* = -19.70$.

In order to estimate the maximum amount of uncertainty induced by our “statistical” K–correction scheme, we have also computed the two parameters of the luminosity function with the extreme assumptions that all galaxies are either ellipticals or spirals. The best fit parameters obtained in these way are $\alpha = -1.15$, $M_{b_J}^* = -19.87$ and $\alpha = -1.05$, $M_{b_J}^* = -19.56$, respectively.

Figure 5 shows the luminosity function (with arbitrary normalization) obtained for our sample: the solid line corresponds to the Schechter functional form obtained through the STY method, while the solid circles have been determined through a modified version of the non–parametric C–Method (Lynden–Bell 1971). The 68% and 95% confidence ellipses for α and $M_{b_J}^*$ are shown in the inset, where the cross represents the best estimate obtained by Loveday et al. (1992) from their survey of galaxies with $b_J \leq 17.15$.

The excellent agreement between our best fit parameters and those obtained by Loveday et al. shows that, up to a depth of $\simeq 600 h^{-1} Mpc$, the

Fig.5: Galaxy luminosity function: solid circles have been computed using a modified version of the C-method and the normalization is arbitrary.

overall shape of the luminosity function is well described by a Schechter function with a faint-end slope $\alpha \sim -1.1$ up to $M = -16$.

Figure 6 shows the 68% and 95% confidence ellipses for α and $M_{b,J}^*$ for galaxies with and without detectable emission lines. The allowed contours are clearly separated at a very high significance level: the galaxies with emission lines have steeper power law slopes and fainter $M_{b,J}^*$. This result is much stronger and qualitatively consistent with the differences in the parameters of the luminosity functions of early- and late-type galaxies found by Loveday et al. (1992).

5. Star Forming Galaxies

The OII doublet, $\lambda 3727$, is the most useful star formation tracer in the blue since this line can serve, although with larger uncertainty, as a substitute of $H\alpha$ (Kennicutt 1992). If R is the number of newly formed stars in solar masses per year, EW the equivalent width of the OII line (in Å), and L the galaxy blue luminosity in solar units, then

$$R = 7 \times 10^{-12} L \times EW \times h^{-2}$$

Fig.6: 68% and 95% confidence ellipses for α and $M_{b,J}^*$ for galaxies with and without detectable emission lines.

(Kennicutt 1992).

The measurement of the equivalent width of the detected emission lines is still incomplete; the following results refer to $\sim 1/3$ of the presently available sample. Moreover a statistical analysis of the emission line properties is further complicated by the fact that the line detectability varies as a function of the spectrum signal to noise ratio. To overcome this problem we have proceeded as follows.

Simple considerations on the line detectability allows to correlate the signal to noise in the line $(S/N)_l$ to its observed equivalent width EW and the signal to noise in the adjacent continuum $(S/N)_c$ as:

$$(S/N)_l \simeq 0.4 \times EW \times (S/N)_c$$

The signal to noise ratio for the continuum has been evaluated for all the objects showing emission lines in two regions of $100 \text{ \AA}$ each at the blue and red sides of the OII line, and the resulting data are shown in Figure 7. The relation corresponding to $(S/N)_l = 10$ is also shown. This relation clearly defines an

Fig.7: Distribution of the observed OII equivalent widths versus signal to noise ratio of the adjacent continuum. The solid line represents a signal to noise in the line $(S/N)_l = 10$.

envelope below which very few lines have been identified visually and measured. We therefore assume that $(S/N)_l \sim 10$ is our detection limit. For galaxies not showing emission lines, $(S/N)_c$ has been evaluated at the expected position of the OII line on the basis of the absorption lines redshift. This allows to estimate the upper limit of the OII equivalent width through the above relation.

We have, therefore, equivalent widths (or corresponding upper limits) of OII $\lambda 3727$ for the whole sample, allowing detailed statistical analysis from $z = 0$ to $z \simeq 0.3$.

The first relevant question we asked is : is there any systematic dependence of OII equivalent width (which is proportional to the number of new stars per unit luminosity) with galaxy luminosity and/or redshift (which is roughly proportional to the lookback time at low z)?

Application of a bivariate regression analysis (ASURV package, Isobe &

Feigelson 1986) to our data (both detections and upper limits) shows that both these correlations are present but with different levels of significance. While the correlation of EW with absolute magnitude (EW decreasing with increasing luminosity) is significant at 4.5σ , the correlation with z (EW increasing with increasing z) is significant at 2.5σ only.

Similar results are obtained by studying the cumulative distribution of EW as a function of M in fixed z bins or as a function of z in fixed M bins. For instance, we have studied the cumulative distributions of OII EW for galaxies of different luminosities in two bins in redshift ($0.13 \leq z \leq 0.185$ and $z \leq 0.09$). With this choice of the redshift bins we are not considering for this analysis the galaxies located in the main peak of the density distribution, where the fraction of emission lines is significantly lower than the average. In both redshift bins faint galaxies have larger OII equivalent widths than bright ones. Part of this effect might be induced by a different mixture of morphological types as a function of absolute magnitude, with the fraction of spirals increasing toward the faint end of the luminosity function; however, the fact that this correlation is significant also for galaxies brighter than $M_{b_J} = -18.8$, where the ratio of spirals to ellipticals is approximately constant, makes us believe that the correlation is true also within a given morphological type.

The dependence of EW versus redshift is more difficult to study for two reasons. First, the redshift interval covered by our data is not very large; second, as already mentioned, the redshift distribution shows several peaks, where a lack of emission line galaxies is found (see Figure 3b). Preliminary results seem to suggest that for a fixed magnitude interval the correlation between OII EW and redshift is not very significant. This is in agreement with the results of the bivariate regression analysis mentioned above. These correlations of OII equivalent width with redshift and/or luminosity will be discussed in greater detail when the whole data set will be available.

Acknowledgements

This work, based on data collected at the European Southern Observatory, has been partially supported through NATO Grant CRG 920150 and EEC Contract ERB-CHRX-CT92-0033.

References

- Broadhurst, T.J., Ellis, R.S., Shanks, T., 1988, MNRAS 235 827
- Broadhurst, T.J., Ellis, R.S., Koo, D.C., Szalay, A.S., 1990, Nature 343 726
- Geller, M.J., Huchra, J.P., 1989, Science 246, 897

- Isobe, T., Feigelson, E.D., 1986, Ap.J. 31, 209
- Heydon-Dumbleton, N.H., Collins, C.A., MacGillivray, H.T., 1988, in *Large-Scale Structures in the Universe*, ed. W. Seitter, H.W. Duerbeck and M. Tacke, Springer-Verlag, p. 71
- Kennicutt, R.C., 1992, Ap.J. 388, 310
- Loveday, J., Peterson, B.A., Efstathiou, G., Maddox, S.J., 1992, Ap.J. 390 338
- Lynden-Bell, D., 1971, MNRAS 155 95
- Sandage, A., Tamman, G., Yahil, A., 1979, Ap.J. 232 252
- Shanks, T., Stevenson, P.R.F., Fong, R., MacGillivray, H.T., 1984, MNRAS 206 767
- Tonry, J., Davis, M., 1979, Ap.J. 84 1511